
Restricted marine
Marginal marine to inner neritic
Outer neritic
Oceanic
Thickness (m)
Stage
E Foram Zones
P Foram Zones
NP Nannozone
CP Nannozone
CN Nannozone
Polarity/Chron
Lithology
Goniodomids
Deflandrea cpx
Phthanoperidinium spp.
Wetzelielloids
Areoligeraceans
Cleistosphaeridium cpx
Cribro-Cordo cpx
Spiniferites pseudofurcatus
Thalassiphora spp.
Protoperidinioids
Enneadocysta-Areosph. cpx
Operculodinium spp.
Cerebrocysta cpx
Spiniferites ramosus cpx
Impagidinium cpx
Nematosphaeropsis spp.
Lingulodinium spp. (no specific eco-group)
Priabonian
Bartonian
E14
E13
E12
E10-11
P15
P14
P13
P12
NP19-20
NP18
NP17
NP16
CP15b
CP15a
CP14b
CP14a
CNE18
CNE17
CNE16
CNE15
CNE14
C16r
C17n.1n
C17n.1r
C17n.2n
C17n.2r
C17n.3n
C17r
C18n.1n
C18n.1r
C18n.2n
C18r
Canaletto
Tiepolo
Tintoretto
Tiziano
Canova
Palladio
Giorgione
Mantegna
GSSP
Sapropelic interval
MECO
(%)